

Constructed Wetlands to Reduce Nutrients From Cropland Runoff: **IMPLICATIONS FOR URBAN STORMWATER**




INAFSM | September 6, 2018

Al Walus

Dave Buckley, PE, CFM, CPSWQ

Mahsa Izadmehr, PhD Candidate





PROJECT OVERVIEW:


Using the Farmable Wetland Program under the U.S. Department of Agriculture's Conservation Reserve Program (CP-39):

The Wetlands Initiative (TWI) has successfully facilitated the design and construction of a wetland treatment located on a private farm in north central Illinois.

The project included two monitoring locations, at the inflow and outflow, allowing for the measurement of nutrient concentrations throughout the growing seasons and periods of dormancy.

Why are we doing this?

- Nutrient runoff is primarily responsible for the annual "dead zone" in the Gulf of Mexico and large algal blooms in parts of the Great Lakes.
- Row-crop agriculture is the biggest source of nutrients.
- Gulf of Mexico Hypoxia Action Plan
 - Requires all watershed states to develop a plan to reduce their nutrients.
- Illinois Nutrient Loss Reduction Strategy
 - Address point-source, urban runoff, and agricultural runoff



Illinois Nutrient Loss Reduction Strategy

- Using strategies from other states, Illinois sought input from major agricultural commodity organizations to support the strategies identified.
 - Illinois Farm Bureau
 - Fertilizer and Chemical Association
 - Corn Growers Association



ILLINOIS
NUTRIENT LOSS
REDUCTION STRATEGY

Improving our
water resources
with collaboration
and innovation

Nutrient pollution is a major threat to water quality in Illinois. Over the decades, state and local efforts to control nutrients have yielded positive results, but new strategies are needed to improve the effectiveness of existing water quality programs and secure the long-term health of water bodies in Illinois and throughout the Mississippi River Basin.

What is nutrient pollution?

Plants and animals need nitrogen and phosphorus to survive. But when too much of either is carried in runoff from city streets and farm fields or flows out of wastewater treatment plants, it can fuel algal blooms that decrease oxygen needed by aquatic plants and animals. In the Gulf of Mexico, nutrients washed down by the Mississippi River have created a 'dead zone' that covers thousands of square miles. Algal blooms also lower property values, hinder recreation, and threaten public health. In addition, nutrient pollution can degrade drinking water quality and require utilities to install costly treatment equipment.



What is Illinois doing to address the problem?

To help protect local streams and the Gulf, Illinois and 11 other states in the Mississippi River Basin have pledged to develop strategies to reduce the nutrient loads leaving their borders.

These strategies are part

of a national plan developed by the Mississippi River, Gulf of Mexico Watershed Nutrient Task Force to reduce the size of the Gulf of Mexico hypoxic zone.

The Illinois Nutrient Loss Reduction Strategy builds on existing efforts by state and local governments, as well as non-profits and industry, to protect and restore Illinois waterways.

Key Strategy Components

1. Extends ongoing regulatory and voluntary efforts
2. Identifies priority watersheds for nutrient loss reduction efforts
3. Establishes the Nutrient Monitoring Council to coordinate water quality monitoring efforts by government agencies, universities, non-profits, and industry
4. Creates the Nutrient Science Advisory Committee to develop numeric nutrient criteria for Illinois waters
5. Forms the Agricultural Water Quality Partnership Forum to oversee outreach and education efforts
6. Establishes the Urban Stormwater Working Group to coordinate and improve stormwater programs and education
7. Lays out strategies for improving collaboration among government, non-profits, and industry
8. Defines a process for regular review and revision

Illinois Nutrient Loss Reduction Strategy



What is Illinois doing to address the problem?

To help protect local streams and the Gulf, Illinois and 11 other states in the Mississippi River Basin have pledged to develop strategies to reduce the nutrient loads leaving their borders.

These strategies are part of a national plan developed by the Mississippi River, Gulf of Mexico Watershed Nutrient Task Force to reduce the size of the Gulf of Mexico hypoxic zone.

The Illinois Nutrient Loss Reduction Strategy builds on existing efforts by state and local governments, as well as non-profits and industry, to protect and restore Illinois waterways.

Illinois Nutrient Loss Reduction Strategy: Key Strategy Components

- 1. Extends ongoing regulatory and voluntary efforts**
- 2. Identifies priority watersheds for nutrient loss reduction efforts**
- 3. Establishes the Nutrient Monitoring Council to coordinate water quality monitoring efforts by government agencies, universities, non-profits, and industry**
- 4. Creates the Nutrient Science Advisory Committee to develop numeric nutrient criteria for Illinois waters**


- 5. Forms the Agricultural Water Quality Partnership Forum to oversee outreach and education efforts**
- 6. Establishes the Urban Stormwater Working Group to coordinate and improve stormwater programs and education**
- 7. Lays out strategies for improving collaboration among government, non-profits, and industry**
- 8. Defines a process for regular review and revision**

Illinois Council on Best Management Practices “What’s your Strategy”

- Illinois Council’s website is the one-stop hub
- The Council focuses on a **system** of practices, with no single best management practice



IL COUNCIL ON BMPs
www.illinoiscbmp.org



BEST MANAGEMENT PRACTICES

Illinois' Nutrient Loss Reduction Strategy (NLRs) characterizes best management practices (BMPs) for agriculture into three categories: In field, Edge-of-field, and Land Use Changes.

The NLRs included BMPs that have been proven to reduce nutrient losses in peer-reviewed, published research. **Those practices are noted here with an ***. This website also includes information about additional BMPs that can be used by farmers here in Illinois, many of which are being tested in ongoing research.

You can find information about Illinois' Demonstration Farms, where many of these practices are being researched and showcased, [here](#).

IN FIELD:


- [Nitrogen management](#) *
- [Cover crops](#) *
- [Reduced tillage](#) *

EDGE-OF-FIELD:

- [Buffers](#) *
- [Saturated buffers](#)
- [Woodchip Bioreactors](#) *
- [Wetlands](#) *
- [Drainage Water Management](#)
- [Streambank Stabilization](#)

LAND USE CHANGES:

- [Perennial/energy crops](#) *



BEST MANAGEMENT PRACTICES

Illinois' Nutrient Loss Reduction Strategy (NLRs) characterizes best management practices (BMPs) for agriculture into three categories: In field, Edge-of-field, and Land Use Changes.

The NLRs included BMPs that have been proven to reduce nutrient losses in peer-reviewed, published research. **Those practices are noted here with an ***. This website also includes information about additional BMPs that can be used by farmers here in Illinois, many of which are being tested in ongoing research.

You can find information about Illinois' Demonstration Farms, where many of these practices are being researched and showcased, [here](#).

IN FIELD:


- [Nitrogen management *](#)
- [Cover crops *](#)
- [Reduced tillage *](#)

EDGE-OF-FIELD:

- [Buffers *](#)
- [Saturated buffers](#)
- [Woodchip Bioreactors *](#)
- [Wetlands *](#)
- [Drainage Water Management](#)
- [Streambank Stabilization](#)

LAND USE CHANGES:

- [Perennial/energy crops *](#)



WHAT'S YOUR STRATEGY?

The Illinois Nutrient Loss Reduction Strategy: Illinois farmers across the state share their conservation stories



[BMPs](#) > [Partners](#) > [About](#) > [Calendar](#)

Expo shows wetland built for nutrient loss reduction



Farmer/Landowner: Thacker Farms


Organizations Involved: The Wetlands Initiative, IL LICA, NRCS, FSA, IL Corn Growers Association

Programs utilized: Conservation Reserve Program (CRP) cost-share

Best Management Practice(s):

- [Constructed Wetland](#)

Description: Bureau County's first constructed wetland for cropland tile drainage treatment was installed at Thacker Farms during a three-day field expo held August 4-6, 2015. This practice is designed to reduce nutrient losses by siting a small wetland along a stream or drainage way where it can intercept tile line(s) and naturally remove excess nutrients. Nearly 100 people from



WHAT'S YOUR STRATEGY?


The Illinois Nutrient Loss Reduction Strategy: Illinois farmers across the state share their conservation stories



Farmer/Landowner: Thacker Farms



where it can intercept the line(s) and naturally remove excess nutrients. Nearly 100 people from



Description: Bureau County's first constructed wetland for cropland tile drainage treatment was installed at Thacker Farms during a three-day field expo held August 4-6, 2015. This practice is designed to reduce nutrient losses by siting a small wetland along a stream or drainage way where it can intercept tile line(s) and naturally remove excess nutrients. Nearly 100 people from 13 counties attended the expo to learn about the wetland and other conservation practices hands-on.





Search 

ABOUT US

EVENTS

NEWS

BEST MANAGEMENT PRACTICES

NUTRIENT LOSS REDUCTION
STRATEGY

WHAT ILLINOIS FARMERS ARE
DOING

WATER TESTING

PRIORITY WATERSHEDS

AGRICULTURE & CONSERVATION
ORGANIZATIONS

CONSERVATION PROGRAMS &
TOOLS

<http://illinoiscbmp.org/> EDUCATION

CONSTRUCTED WETLANDS

A wetland is a marsh-type area with saturated soils and water-loving plants. Wetlands can be constructed for the purpose of removing nutrients because they filter nutrients, chemicals, and sediment from runoff or tile water before water moves off of a farm field and into streams and rivers. Because wetlands slow overland flow and store runoff water, they reduce both soil erosion and flooding downstream. Many wetlands release water slowly into the ground which recharges groundwater supplies.


Wetlands **provide habitat for waterfowl and many other species** of wildlife, as well as add beauty and value to a farm. Wetlands can be built or enhanced by installing practices such as dikes to manage water levels.

Wetlands targeted for water quality benefits show great potential for nitrate-N reduction. Wetland costs include design, construction, buffer seeding, maintenance and land acquisition. In addition to water quality benefits, these wetlands provide other benefits such as improved aesthetics and habitat.

Nature's Kidneys: [The Illinois Water Resources Center](#) discuss how constructed wetlands filter nitrogen and benefit wildlife in the video below.



RECEIVE OUR E-NEWSLETTER



Current Conditions:

We need **both** fertilizer and drainage for productive farming.

One of the least expensive ways to address nutrient runoff is through the rate and timing of fertilizer applications.

However, even the most careful farmer can't avoid some nutrient loss. This is largely due to the drain tile system.

The drain tile has been a critical aspect to farming since the mid-19th century responsible for making planting and harvesting more consistent and reliable from year to year.



The drain tile acts as a transport vehicle, allowing field drainage of excess water to carry nutrients with it, including nitrates.


Cropland Treatment Practices – BMPs

- We can achieve significant nitrate reduction by treating nutrients leaving the field through drain tiles with:

- Vegetated Buffers
- Bioreactors
- Constructed Wetlands


Constructed Wetlands

- ❖ Specifically located and designed for a particular drainage area for the purpose of intercepting drain tile drainage to reduce nutrients before reaching a receiving waterway.
- ❖ Optimize the natural process to remove nutrients.



“Working wetlands” are one of the most promising practices for reducing nutrient loss.

Comparison of nitrogen removal cost-effectiveness for select agricultural practices
(estimated average annual cost in \$/pound of nitrogen removed)



Constructed Wetland




- Densely vegetative marsh versus open water
- Vegetation is critical to slow water down while providing substrate for working microbes
- 50 year functionality with very low maintenance
- Provides environmental benefits
 - Pollinator habitat
 - Wildlife habitat
 - Carbon sequestration

Buy-In and Cost Share Strategy

- **The Wetlands Initiative works with farmers (1 on 1) to promote interest.**
 - TWI is a non-profit organization dedicated to restoring the wetland resources of the Midwest.
 - Land owner confidence that the practice will work.
 - Local buy-in, trusted farm leaders.
 - Minimizing impacts to farming operations.
 - Implemented in often low producing areas of the farm.
- **Not simply building a few wetlands and assume other farmers will copy and take action.**
 - TWI is spreading the practice within the real-life economics of the working Farm Belt.
 - TWI wants to prove this type of on-the-ground conservation is not some little boutique thing but a normal part of the working farm-belt landscape just like nutrient management, grassed waterways or drainage ditches.
- **Federal cost share programs → Farm Service Agency**
 - Offset the cost for this practice while reducing investment in less-profitable land.
 - EQIP – Environmental Quality Incentives Program
 - CRP – Conservation Reserve Program
 - Is the project eligible
 - NRCS – must approve the design.



Thacker Farms, West Bureau Creek near Princeton IL





1 POSITION

- Intercept tile drainage before outlet ditch or stream
- Capture high nutrient loads
- Locate in watershed headwater areas
- Marginal or unprofitable land

2 SIZE

- Key to nutrient removal
- Allow adequate residence time
- Treatment area is 0.5-5.0% of the drainage area
- Treatment area is 12" above to 24" below permanent pool

3 DEPTH

- Marsh wetland (aka shallow "pond")
- At least 50% of the permanent (normal) pool is 12" or less
- Anything greater than 24" in depth doesn't count towards the ratio or treatment area


Design

NRCS Criteria

- HMS Hydrologic Modeling
- SCS Methodology
 - 25-yr, 24-hr
 - Max velocity = 1.5 ft/sec
 - 72-hr draw down; 10yr, 24-hr storm


Design

- Located adjacent to creek
- Inlet and outlet structures
- 40 acres of tributary area
- Treatment area is 0.5 acres
- Small berms to increase flow path



Planting Plan

- Total footprint is 4.3 acres
- Wetland area
- Different seed mixes each zone



Construction



In an effort to increase public awareness and education, TWI partnered with the IL chapter of the Land Improvement Contractors of America.


The wetland was built as part of ILICA's conservation expo that was held Aug 4-6th. The construction was between the 3rd- 8th.



7000 Cubic Yards

Compacted Clay
Liner






Embedded Video from 2015 Conservation Expo (Ohio, IL)
has been removed from this presentation due to size limitations

Performance Monitoring by UIC Dept. of Civil and Materials Engineering

Sampling at Inflow and Outflow




Wetland water level controlled by outlet weir




Flow calculated from known geometry and measured stage using the Francis weir equation

$$Q = 1.838LH^{3/2}$$





Wetland Hydraulics: Tracer Study

- Bromide (Br^-) tracer injected for 6 hrs at inlet
 - Sampling at wetland cell mid-point and the outlet control structure
 - Conductivity at 5 min intervals, Br^- at 1 hr intervals
 - Overall recovery was $\sim 90\%$
- 1D Transport with Inflow and Storage (OTIS) model
 - No substantial short-circuiting
 - Substantial dispersion
 - Clear peak tailing in the outlet tracer
 - $\text{HRT} = 17.5 \pm 6.7 \text{ hr}$




P removal > N removal (on a % basis)



Overall, removal averaged 22% for SRP (Soluble Reactive Phosphorus), 10.6% for nitrate and 10.3% for TIN (Total Inorganic Nitrogen)

Cumulative N and P mass removal by the system



Based on the measured flowrates and inlet/outlet nutrient concentrations, the cumulative N and P removal was determined using a mass balance approach.

Approximately 120 kg of SRP and 1200 kg (1.3 tons) of NO3-N (Nitrate as Nitrogen)

Why was N removal efficiency low?


N removal is primarily through **denitrification of NO_3 to produce N_2**

Denitrification requires the presence of three (3) components simultaneously:

1. Electron Donors (simple organic compounds from breakdown of organic material)
2. Electron Acceptors (i.e. NO_3 from fertilizers and nitrification of NH_3)
NOT AN ISSUE HERE!
3. Competent microbes to carry out the process (i.e. denitrifying bacteria)

Thus, possible reasons for lack of/low N removal include:

1. Lack of Electron Donors (not much organic material at start)
2. No or low levels of denitrifying bacteria present at start
3. Hydraulic overloading (Electron Acceptors overwhelms available Electron Donors)



Why was N removal low?

Investigating these one by one:


1. Lack of Electron Donors (plenty of Electron Acceptors!)
 - Although the sediment is **~4% Organic Material**, this Organic Material may not be highly biodegradable and thus may not produce sufficient amounts of Electron Donors to match the Electron Acceptor load
 - Further monitoring of Organic Material levels will help determine whether they increase from wetland growth and development
2. Lack of competent microbial community structure
 - It is likely that denitrifying bacteria need time to adapt to the wetland conditions with abundant NO_3 levels
 - Further monitoring of N removal and microbial community structure analysis via 16S RNA sequencing is ongoing
3. Overloading (Electron Acceptors overwhelms available Electron Donors)
 - It is possible that the higher flowrates resulted in NO_3 overloading
 - 17.5 hr HRT in the tracer study was lower than we expected, resulting in less time for denitrification to occur
 - We expect possibly longer HRT now that the weir depth is fixed and wetland plants established

Wetland development: From planting to operation

Dec 2015



March 2016



Jun 2016




Aug 2016



Increased incorporation of new labile Organic Material into sediments from wetland plant growth

This will result in increased N removal efficiency (more denitrification) from our constructed “kidney”

Constructed Wetlands to Reduce Nutrients From Cropland Runoff: **IMPLICATIONS FOR URBAN STORMWATER**



Construction photos and thanks
to all involved:

Conservation Expo 2015



August 4 – 6, 2015

